

UNC CFAR Social and Behavioral Science Research Core SABI Database

INSTRUMENT TITLE: Mausbach Self-Efficacy for Condom Use

SOURCE ARTICLE: Mausbach, B. T., Semple, S. J., Strathdee, S. A., Zians, J., & Patterson, T. L. (2007). Effectiveness of a behavioral intervention for increasing safer sex behaviors in HIV-positive MSM methamphetamine users: Results from the EDGE study. *Drug and Alcohol Dependence*, 87(2-3), 249.

RESPONSE OPTIONS AND SCORING: Responses for each item ranged from 0="strongly disagree" to 3="strongly agree". Responses to these nine items were summed to create an overall self-efficacy score (range = 0–27).

SURVEY ITEMS:

- 1. I can use a condom properly
- 2. I can use a condom every time I have penetrative sex
- 3. I can have condoms available every time I have penetrative sex
- 4. I can use a condom in any situation
- 5. I can interrupt sex to use a condom
- 6. I can use a condom without any instruction
- 7. I can use a condom for penetrative sex while under the influence of drugs or alcohol
- 8. I can delay penetrative sex if a condom is not available
- 9. I can use a condom for penetrative sex even when I am very sexually aroused

TERMS OF USE:

Individuals may use this information for research or educational purposes <u>only</u> and may not use this information for commercial purposes. When using this instrument, please cite:

Mausbach, B. T., Semple, S. J., Strathdee, S. A., Zians, J., & Patterson, T. L. (2007). Effectiveness of a behavioral intervention for increasing safer sex behaviors in HIV-positive MSM methamphetamine users: Results from the EDGE study. Drug and Alcohol Dependence, 87(2-3), 249.

When presenting results using any survey information you obtained from the SABI, please acknowledge the University of North Carolina at Chapel Hill Center for AIDS Research (CFAR), an NIH funded program P30 AI50410.